

HMR and RMMR Fact sheet for medical specialists



Medicines are the most common form of treatment used in healthcare. Patients require more medicines as they age, and their medical needs become more complex.¹ Almost one in two Australians (47%) have a chronic medical condition, with 80% of people aged 65 years and over having one or more chronic conditions.² Medication-related problems are common, especially during transitions of care.³ Collaborative medication reviews support optimal medication use and improve patient health outcomes.⁴



Over 90% of patients have at least one medication-related problem post-discharge from hospital



Over 95% of residents living in aged care homes have at least one medication-related problem; most have three problems



Over half of residents in aged care homes are prescribed medicines that are considered potentially inappropriate in older people



40% - 50% of people living in aged care homes are on medicines that have the potential to cause sedation or confusion



20% of people living in aged care homes are on antipsychotics; more than half use the medicine for too long



Up to one-third of people living in aged care are taking benzodiazepines; more than half use the medicine for too long

Quality Use of Medicines and Medication Safety is Australia's 10th National Health Priority

Medication safety and quality use of medicines in Australia

There are 250,000 hospital admissions each year due to medication-related problems. Half of this harm is considered to be preventable.¹ Medication-related problems are responsible for 400,000 emergency department presentations each year, and in the last 6 months 1.2 million Australians have experienced an adverse medication event.¹

How can Home Medicines Reviews (HMRs) and Residential Medication Management Reviews (RMMRs) support better medication management and reduce medication-related harm?

- On average, four medication-related problems are detected for each person who has a HMR, the majority of which can be resolved.¹
- Medication reviews conducted by pharmacists can reduce medication-related problems and improve adherence.⁷
- HMRs are associated with a reduction in hospitalisation rates for older people living in the community at high risk of medication-related hospital admissions (e.g. those with heart failure taking heart failure medicines⁸ and those taking warfarin⁹).
- RMMRs are effective in identifying medication-related problems in aged care homes (on average 2.7-3.9 medication-related problems per resident¹⁰).
- Targeted activities in aged care homes including audit and feedback, education and medication reviews reduce inappropriate use of psychotropic medicines, opioids, and antimicrobials.⁵



Who will benefit from a comprehensive medication management review?

A comprehensive medication management review (HMR or RMMR) could benefit a person who is at risk of medication-related harm due to:

- multiple chronic conditions or comorbidities
- age
- social circumstances
- characteristics of their medicine
- complexity of their medication regimen
- limited knowledge and skills to use their medicines effectively and safely.

Table 1 lists examples of risk criteria to identify patients who may benefit from an HMR or RMMR. These are not mandatory criteria nor a comprehensive list but are intended as a guide. Number of medicines (e.g. 5 or more medicines) is not a valid indicator for risk of medication-related harm.

The referral to an accredited pharmacist should include specific clinical information relating to these risk factors and the patient's clinical status.

Pharmacists may conduct up to two follow-ups within 9 months of the initial HMR or RMMR, with a brief report to the referring medical practitioner.


Post-discharge medication reviews can be triggered by a hospital doctor, with support from a hospital pharmacist, where available. They have an additional focus on continuity of care, medication reconciliation, and immediate patient safety and action.

There is no MBS Item number for medical specialists for HMRs or RMMRs.

Table 1. Risk factors suggesting a patient may benefit from a comprehensive medication management review

Goals of medication therapy not reached or maintained, including suboptimal response to medication
Chronic medical condition associated with a high risk of unplanned hospital admission (e.g. chronic obstructive pulmonary disease, heart failure, chronic pain)
Three or more chronic medical conditions
Recent discharge from hospital or frequent unplanned hospital admissions
Recent admission to a residential aged care facility
Significant changes to medication regimen, including newly prescribed medicines
High-risk medicines requiring close monitoring for adverse effects and/or efficacy (e.g. opioids, psychotropic medicines, insulin, anticoagulants, antibiotics, NSAIDs, anticholinergics)
Functional issues that increase the risk of harm and/or reduce the chance of benefit from medicine use (e.g. frailty, frequent falls, cognitive impairment, swallowing difficulty, renal or hepatic impairment)
Symptoms suggestive of any adverse drug reaction
Prescribing cascade (e.g. one medicine to treat an adverse effect of another)
Problems managing medicine-related therapeutic devices (e.g. inhalers, subcutaneous injections, eye drops, transdermal patches)
Difficulty understanding and following medication regimen
Language, literacy or cultural difficulties

MEDICATION REVIEW CYCLE OF CARE



References:

1. Pharmaceutical Society of Australia 2019. Medicine Safety: Take Care. Canberra: Pharmaceutical Society of Australia; 2019.
2. Australian Bureau of Statistics. National Health Survey: First Results, 2017-18. 2018. At: www.abs.gov.au/ausstats/abs@nsf/mf/4364.0.55.001
3. Wheeler A, Scatill S, Hopcroft D, et al. Reducing medication errors at transitions of care is everyone's business. Aust Prescr 2018;41(3):73-7.
4. Castelino RL, Bajorek BV, Chen TF. Retrospective evaluation of home medicines review by pharmacists in older Australian patients using the medication appropriateness index. Ann Pharmacother 2010;44(12):1922-9.
5. Pharmaceutical Society of Australia. Medicine safety: aged care. Canberra: Pharmaceutical Society of Australia; 2020.
6. Pharmacy Programs Administrator. Program rules: Home medicines review. 2020. At: www.ppaonline.com.au/wp-content/uploads/2019/01/HMR-Program-Rules.pdf
7. Jokanovic N, Tan ECK, van den Bosch D, et al. Clinical medication review in Australia: A systematic review. Res Social Adm Pharm 2016;12:384-418.
8. Roughead EE, Barratt JD, Ramsay E, et al. The effectiveness of collaborative medicine reviews in delaying time to next hospitalization for patients with heart failure in the practice setting. Circ Heart Fail 2009;2:424-8.
9. Roughead EE, Barratt JD, Ramsay E, et al. Collaborative home medicines review delays time to next hospitalization for warfarin associated bleeding in Australian war veterans. J Clin Pharm Ther 2011;36(1):27-32.
10. Chen EYH, Wang KN, Slugggett JK, et al. Process, impact and outcomes of medication review in Australian residential aged care facilities: A systematic review. Australas J Ageing. 2019;38 Suppl 2:9-25.